

woman chooses to reveal such fears, worries, and concerns to the doctor.³

There is ample evidence of patients' frequent dissatisfaction with doctors' communication skills.⁴ Despite increasingly liberal provision of information, many patients still want to know more than they are told,⁵ whereas others do not want to participate in decision making; they need absolute, uncritical confidence in their doctors' skills. A skilful doctor will achieve the correct balance between autonomy and paternalism for each patient. To determine the appropriate balance it is useful to have a framework to help identify the necessary tasks and skills.

Such a framework is provided by the "three function" model of the consultation developed by Bird and Cohen-Cole⁶: gathering data to understand the patient; developing rapport and responding to the patient's emotions (to enable the patient to feel understood); and patient education and behaviour management. These functions relate to the three purposes and effects of communication: informative (to exchange information); promotive (to bring about action); and evocative (to arouse certain feelings)⁷ and also to the three domains (cognitive, affective, and psychomotor) widely used by educationalists to categorise educational objectives. Each function has specific objectives and demands specific explicit skills of the doctor if they are to be achieved. Fortunately, considerable evidence now exists that such skills can be successfully acquired.⁸

However, no single model can fully convey the complexity of the doctor-patient relationship and the three function model needs to be expanded by drawing on other concepts of the consultation. These include the idea of the consultation as a "meeting between experts"⁹; the patient-centred clinical method described by Stewart et al¹⁰; the problem based approach¹¹; the stages of motivational interviewing described by Miller and Rollnick¹²; and the family systems approach, which emphasises the importance of taking into account the patient's family and social networks.¹³

Finally, the individuality of the professional cannot be ignored. All sorts of factors, some based on the pro-

fessional's own life experiences, can both consciously and unconsciously influence his or her behaviour and decisions. This has led to the notion of the doctor as a drug,¹⁴ with both powerful effects and side effects. When listening and talking to patients, professionals need to be aware not only of the words they use, both to discover and convey information, but also their own feelings and how to cope with them.

Thus a recommendation to change terminology, while laudable in its intentions, may not be enough to alleviate mothers' dissatisfaction with the care that they receive. The risk is that mere use of the "correct" terminology, with no attention paid to the wider aspects of a consultation, could lead to professional complacency.

Paul Freeling *Emeritus professor of general practice*

St George's Hospital Medical School, London SW17 0RE

Linda Gask *Senior lecturer in psychiatry*

National Primary Care Research and Development Centre, University of Manchester, Manchester M13 9PS

- 1 Hutchon DJR, Cooper S. Terminology for early pregnancy loss must be changed. *BMJ* 1998;317:1081.
- 2 Roberts AH. Managing miscarriage: the management of the emotional sequelae of miscarriage in training practices in the West of Scotland. *Fam Pract* 1991;8:117-20.
- 3 Goldberg DP, Jenkins L, Millar T, Faragher EB. The ability of trainee general practitioners to identify psychological distress among their patients. *Psychol Med* 1993;23:185-93.
- 4 Audit Commission. *What seems to be the matter? Communication between hospitals and patients*. London: HMSO, 1993.
- 5 Stott NC, Pill RM. Advise yes. Dictate no. Patients' views on health promotion in the consultation. *Fam Pract* 1990;7:125-31.
- 6 Bird J, Cohen-Cole SA. The three function model of the medical interview: an educational device. *Adv Psychosom Med* 1990;20:65-88.
- 7 Browne K, Freeling P. *The doctor patient relationship*. 2nd ed. Edinburgh: Churchill Livingstone, 1976.
- 8 Simpson M, Buckman R, Stewart M, Maguire P, Lipkin M, Novack D, et al. Doctor-patient communication: the Toronto consensus statement. *BMJ* 1991;303:1385-7.
- 9 Tuckett D, Boulton M, Olson C, Williams A. *Meetings between experts: an approach to sharing ideas in medical consultations*. London: Tavistock, 1985.
- 10 Stewart M, Brown BJ, Weston WA, McWhinney I, McWilliam CL, Freeman TR. *Patient centered medicine: transforming the clinical method*. Thousand Oaks: Sage, 1995.
- 11 Lesser AL. Problem based interviewing in general practice: a model. *Med Educ* 1998;19:299-304.
- 12 Miller WR, Rollnick S. *Motivational interviewing: preparing people to change addictive behaviour*. New York: Guilford Press, 1991.
- 13 McDaniel A, Campbell T, Seaburn D. *Family orientated primary care: a manual for medical providers*. New York: Springer Verlag, 1992.
- 14 Balint M. *The doctor, his patient and the illness*. Pitman: London, 1957.

Fluoroquinolone resistance

Overuse of fluoroquinolones in human and veterinary medicine can breed resistance

Fluoroquinolones are highly active, broad spectrum antibiotics with many uses in both human and veterinary medicine. As with other classes of antibiotic, however, decreased susceptibility or resistance to these agents has developed. Resistance is usually chromosomally mediated so the spread of resistant bacteria contributes to the high numbers of resistant strains reported by some institutions. Although for many infections the changes in susceptibility of the bacteria have not had an immediate clinical impact, highly resistant strains have recently emerged. For some infections and bacterial species the clinical usefulness of fluoroquinolones may be limited if their use is not curtailed.

Genetic and biochemical experiments have shown two enzymes that are the targets of fluoroquinolones.¹ In Gram negative bacteria such as *Escherichia coli*, *Pseudomonas aeruginosa*, and *Neisseria gonorrhoea*, the primary target is DNA gyrase and the secondary target DNA topoisomerase IV; in Gram positive bacteria, such as *Staphylococcus aureus* and *S pneumoniae*, the primary and secondary targets are reversed.² DNA gyrase and topoisomerase IV are each enzymes composed of four subunits (two A and two B) encoded by *gyrA* and *gyrB*, and *parC* and *parE* respectively. Fluoroquinolone resistant isolates usually contain one or more mutations in a small section of *gyrA* or *parC*; mutation

in *gyrB* and *parE* is rare.¹ In bacteria where mutations have given rise to a resistant DNA gyrase mutations then occur in the topoisomerase IV genes (and vice versa for Gram positive bacteria) to give a highly resistant bacterium. In addition there are genes that influence the uptake of the drug into the bacterial cell and efflux pumps that can be overexpressed to enhance excretion of quinolones from the cell. This enhanced efflux in turn causes increased minimum inhibitory concentrations of several drugs, including fluoroquinolones, tetracycline, chloramphenicol, and ampicillin.³⁻⁶ It has been suggested that mutations enhancing efflux occur as a first step, allowing the bacteria to survive such that mutations can accumulate in genes encoding the target proteins. Plasmid mediated, transferable, fluoroquinolone resistance has recently been described⁷; its mechanism of resistance and epidemiology is unknown.

Gram negative bacteria such as *Enterobacteriaceae* spp and *N gonorrhoea* are usually exquisitely susceptible to fluoroquinolones, and repeated exposure to fluoroquinolones has probably allowed the bacteria to acquire and accumulate the mutations that give rise to resistance. Highly resistant *E coli* and *N gonorrhoea* have already been described.⁸⁻⁹ For gut commensals such as *E coli*, frequent exposure to fluoroquinolones during treatment of other infections, such as those of the respiratory tract, may allow the accretion of mutations and emergence of highly resistant bacteria. In susceptible hosts these then give rise to infections. Although decreased susceptibility of gonococci to fluoroquinolones has become widespread, clinical resistance remains rare, occurring only in geographical hot spots such as the Far East. Clearly clonal spread of resistant bacteria is important in the spread of fluoroquinolone resistant gonococci, so efforts to reduce transfer of the bacteria are necessary.

Some bacteria such as *Salmonella typhimurium* and *Campylobacter* species are well known zoonoses. *S typhimurium* with decreased susceptibility to fluoroquinolones and *Campylobacter* resistant to fluoroquinolones have been isolated from animals and retail poultry.¹⁰⁻¹¹ For these reasons the use of fluoroquinolones in veterinary medicine has caused concern, as such strains can infect man. A fluoroquinolone is often the drug of choice for patients who have chronic enteritis, are immunocompromised, or have extraintestinal infection. There is evidence that complicated infections caused by resistant strains of *S typhimurium* are difficult to treat; *S typhi* (not a zoonose) with reduced susceptibility to fluoroquinolones (resistant to nalidixic acid) responded more slowly to fluoroquinolone treatment—and clinical outcome was poorer.¹² Therefore, it has been suggested that nalidixic acid should be used to detect decreases in susceptibility which may herald fluoroquinolone resistance. Many patients (>60%) infected with a nontyphoidal salmonella with decreased susceptibility to fluoroquinolones or a fluoroquinolone resistant *Campylobacter* have had no prior exposure to fluoroquinolone, suggesting that the bacteria must have been ingested. However, it is unclear how many patients have received a fluoroquinolone to treat their infection and whether their clinical outcome is poor. Until the full clinical relevance and risk to human health has been determined the role of fluoroquinolones in veterinary medicine remains controversial.

Another major use of fluoroquinolones is in treating respiratory tract infections, especially with the development of agents to treat pneumococcal infections. Gram positive bacteria are inherently less susceptible to fluoroquinolones than many Gram negative bacteria, so bacteria such as *Streptococcus pneumoniae* typically require only one, sometimes two, mutations in the genes encoding the target proteins to become clinically resistant. Therefore fluoroquinolone resistance is more likely to occur. For *Staphylococcus aureus* clonal spread of these bacteria in institutions has always been a problem and now fluoroquinolone resistance has been added to methicillin resistance. Good infection control will reduce the transfer and prevalence of these strains.

Fluoroquinolones have also been used as second line agents to treat multidrug resistant tuberculosis. However, mycobacteria have also been able to acquire mutations in the genes encoding the target protein.¹³ Such strains are prevalent only in areas where fluoroquinolones have been widely used, and preventing the spread of *Mycobacterium tuberculosis* will reduce the numbers of resistant bacteria isolated.

As with other antibacterial agents, initial optimism about fluoroquinolones has been tempered by the development of resistant strains. The species that have become problematic are typically those that are inherently less susceptible to this class of agents. There has been misuse and abuse, so to retain the excellent activity of this class of antibiotics and reduce the development of resistant strains and their spread fluoroquinolones should be used prudently and only where there is a clinical need. For several infections a fluoroquinolone is not necessary as a first line agent.

Laura J V Piddock *Head*

Antimicrobial Agents Research Group, Division of Immunity and Infection, University of Birmingham, Birmingham B15 2TT (l.j.v.piddock@bham.ac.uk)

- 1 Everett MJ, Piddock LJV. Resistance to quinolones and fluoroquinolones. In: Kahlman B, Dalhoff A, eds. *Handbook of experimental pharmacology: microbial resistance to drugs*. Heidelberg: Springer-Verlag, 1997.
- 2 Pan XS, Fisher LM. Cloning and characterization of the *parC* and *parE* genes of *Streptococcus pneumoniae* encoding DNA Topoisomerase IV—role in fluoroquinolone resistance. *J Bacteriol* 1996;178:4060-9.
- 3 Alekshun MN, Levy SB. Regulation of chromosomally mediated multiple antibiotic resistance: the *mar* regulon. *Antimicrob Agents Chemother* 1997;41:2067-75.
- 4 Miller PF, Sulavik MC. Overlaps and parallels in the regulation of intrinsic multi-antibiotic resistance in *Escherichia coli*. *Mol Microbiol* 1996;21:441-8.
- 5 Poole K, Krebs K, McNally K, Neshat S. Multiple antibiotic resistance in *Pseudomonas aeruginosa*: evidence for involvement of an efflux operon. *J Bacteriol* 1993;175:7363-72.
- 6 Kaatz GW, Seo SM, Ruble CA. Efflux-mediated fluoroquinolone resistance in *Staphylococcus aureus*. *Antimicrob Agents Chemother* 1993;37:1086-94.
- 7 Martinez L, Pascual A, Jacoby GA. Quinolone resistance from a transferable plasmid. *Lancet* 1998;351:797-9.
- 8 Everett MJ, Jin Y-F, Ricci V, Piddock LJV. Contribution of individual mechanisms to fluoroquinolone resistance in 36 *Escherichia coli* isolated from humans and animals. *Antimicrob Agents Chemother* 1996;40:2380-6.
- 9 Ross JDC. Fluoroquinolone resistance in gonorrhoea: how, where and so what. *Int J Sex Transmitted Dis and AIDS* 1998;9:318-22.
- 10 Griggs DJ, Hall MC, Jin YF, Piddock LJV. Quinolone-resistance in veterinary isolates of salmonella. *J Antimicrob Chemother* 1994;33:1173-89.
- 11 Gaunt PN, Piddock LJV. Ciprofloxacin resistant campylobacter in humans: an epidemiological and laboratory study. *J Antimicrob Chemother* 1996;37:747-57.
- 12 Wain J, Hoa NTT, Chinh NT, Vinh H, Everett MJ, Diep TS, et al. Quinolone resistant *Salmonella typhi* in Viet Nam: molecular basis of resistance and clinical response to treatment. *Clin Infect Dis* 1997;25:1404-10.
- 13 Takiff HE, Salazar L, Guerrero C, Philipp W, Huang WM, Kreiswirth B, et al. Cloning and nucleotide-sequence of *Mycobacterium tuberculosis gyrA* and *gyrB* genes and detection of quinolone resistance mutations. *Antimicrob Agents Chemother* 1994;38:773-80.